

Environmental Monitoring and Surveillance of Rodents and Vectors for *Francisella tularensis* Following Outbreaks of Human Tularemia in Georgia

Eka Elashvili,^{1*} Ian Kracalik,^{2,3*} Irma Burjanadze,^{4*} Sophio Datukishvili,⁴ Gvantsa Chanturia,⁴ Nikoloz Tsertsvadze,⁴ Levan Beridze,⁴ Merab Shavishvili,⁴ Archil Dzeladze,⁴ Marina Grdzeldze,⁴ Paata Imnadze,⁴ Andrew Pearson,⁵ and Jason K. Blackburn^{2,3}

Abstract

Tularemia is a re-emerging bacterial zoonosis, broadly distributed across the northern hemisphere. In Georgia, there is a history of human tularemia outbreaks dating back to the 1940s. In response to outbreaks, health officials initiated long-term field surveillance and environmental monitoring. The objective of our study was to obtain information from 57 years of field surveys to identify species that play a role in the occurrence *Francisella tularensis* subsp. *holarctica* in the environment in Georgia. We collected historical data on human outbreaks, field collections, population dynamics of the common vole (*Microtus arvalis*), and conducted surveys on small mammals and vectors from five regions in Georgia during 1956–2012. Bacterial isolation was conducted using standard culturing techniques, and isolation rates for species were obtained for a subset of years. We used a Spearman rank correlation to test for associations between the density of the common vole and isolation rates. From 1956 through 2012, there were four recorded outbreaks of human tularemia (362 cases). A total of 465 bacterial isolates of *F. tularensis* subsp. *holarctica* were obtained from 27 species and environmental samples. The number of isolations was highest in the common vole (*M. arvalis*; 149 isolates; 32%) and *Dermacentor marginatus* ticks (132 isolates; 28%); isolation rates ranged between 0–0.91% and 0–0.47%, respectively. Population dynamics of the common vole were not correlated with the isolation rate. Given the history of tularemia re-emergence in Georgia, continued field surveys and environmental monitoring may provide an early indication of outbreak risk in humans. In conclusion, our findings provide evidence of long-standing foci of *F. tularensis* subsp. *holarctica* that are likely maintained by the common vole–tick cycle.

Key Words: *Francisella tularensis*—Human tularemia—Georgia—*Microtus arvalis*—*Dermacentor marginatus*—Vole–tick cycle.

Introduction

TULAREMIA IS A BACTERIAL ZOOLOGIC DISEASE broadly distributed across the Northern Hemisphere. The causative agent of the disease, *Francisella tularensis*, is thought to circulate in the environment among small mammals (primarily rodents and lagomorphs) with arthropods acting as vectors (Petersen and Schriefer 2005, Rossow et al. 2014).

Recently, tularemia has re-emerged in several European countries (Petersen and Schriefer 2005). Despite recent outbreaks and an increasing incidence in parts of Europe (Hestvik et al. 2015), the ecology of *F. tularensis* is still poorly understood.

In Georgia, there is history of tularemia outbreaks in humans dating back to the 1940s among soldiers at a military garrison in the Samtshke-Javakheti region. In response to this

¹National Food Agency, Tbilisi, Georgia.

²Spatial Epidemiology and Ecology Research Lab, Department of Geography, University of Florida, Gainesville, Florida.

³Emerging Pathogens Institute, University of Florida, Gainesville, Florida.

⁴National Center for Disease Control and Public Health, Tbilisi, Georgia.

⁵University of Maryland, Maryland.

*These authors contributed equally to this work.

outbreak and subsequent outbreaks (Fig. 1), the National Center for Disease Control and Public Health (NCDC) in Tbilisi, Georgia, began monitoring the re-emergence of human tularemia through surveillance of small mammals and vectors and the collection of environmental samples. Genetic characterization of bacterial isolates in Georgia showed all belonged to *F. tularensis* (type B) *holarctica* (Chanturia et al. 2011). A recent outbreak of water-borne tularemia in central Georgia has reiterated the need for continued monitoring and surveillance (Chitadze et al. 2009). The objective of our study was to analyze data from environmental sampling and field studies to trap small mammals and vectors aimed at isolating *F. tularensis* subsp. *holarctica* from several regions of Georgia to better understand the ecology.

Materials and Methods

Field surveys of small mammals and vectors were carried out by the NCDC during the period of 1956–2012. Generally, mammals were trapped with snap traps or Sherman-style live traps, and ticks were collected off of mammals or by flagging (Velijanashvili 1992). The study area (23,837 km²) included five regions in southeastern and eastern Georgia—Samtshke, Javakheti, Shida Kartli, Kvemo Kartli, and Kakheti. Mammals and ticks were identified to species and bacterial isolates were obtained via traditional culturing techniques (World Health Organization 2007). We derived the annual rate (%) of bacterial isolation by species (number of isolates/total number collected); data for bacterial isolation rates were limited




FIG. 1. (A) Total number of *F. tularensis* bacterial isolates collected from field surveys of small mammals and vectors and environmental sampling in Georgia 1956–2012. Rates of bacterial isolation by species were only available during 1963–1985. Primary species included *Microtus arvalis*, *Apodemus sylvaticus*, and *Dermacentor marginatus*. Outbreak years (OB) were recorded in 1956 (20 cases), 1984 (300 cases), 1997 (seven cases), and 2007 (35 cases). (B) Sample diversity of *F. tularensis* from small mammals (black bars), hard ticks (dark grey bars), and fleas/mites (light grey bars).

to the Samtskhe-Javakheti region and to the period from 1963 to 1985.

We plotted, over time, the total number of bacterial isolates obtained, bacterial isolation rates, and the dates of human tularemia outbreaks obtained from historical records at the NCDC (sporadic human cases were not included). Here we define an outbreak as five or more cases in a single community. Density estimates of the common vole (*Microtus arvalis*) (derived from the number of active burrows/hectare) were collected in the Samtskhe-Javakheti region during 1963–1985 (Velijanashvili 1992). To test for correlation between rates of bacterial isolation and common vole density, we used a Spearman rank test (SPSS, IBM, New York, NY).

Results

From 1956 to 2012, a total of 465 (range, 0–51) bacterial isolates of *F. tularensis* subsp. *holarctica* were obtained from field surveys (Fig. 1A). During 1963–1985, samples were obtained from 95,399 *M. arvalis*, 9349 *Apodemus sylvaticus*, 2917 *Mus musculus*, 1428 *Arvicola terrestris*, 3916 from 10 other small mammal species, 17,008 Ixodidae ticks, 800,966 fleas, and gamasid mites. Isolates were obtained from 27 different species and environmental samples (e.g., water, wheat); 194 isolates were obtained from small mammals, 184 from hard ticks, 75 from fleas/mites, and eight from environmental samples (Fig. 1B). Most small mammal isolates were obtained from rodents, although seven isolates were from shrews (Family: Soricidae). Common voles and *Dermacentor marginatus* ticks comprised the greatest number of isolates among small mammals and vectors, 32% (149) and 28% (132), respectively. Rates of isolation during 1963–1985 are shown in Figure 1A. The percentage of isolation (total number collected) range for common voles was 0–0.91 (244–8883) and *D. marginatus* was 0–0.47 (25–1206). Density of the common vole in the Samtskhe-Javakheti region ranged between 22 and 196 animals/hectare (Fig. S1) (Supplementary Data are available at www.liebertonline.com/vbz/). There was no significant correlation between the rate of bacterial isolation and vole population density ($\rho=0.282$, $p=0.19$).

Human outbreaks of tularemia were recorded in 1956, 1984, 1997, and 2007 corresponding to human cases totals of 20, 300, 7, and 35, respectively (Fig. 1A).

Discussion

We obtained data from several thousand small mammals, vectors, and environmental samples collected from 57 years of field surveys in Georgia, resulting in the recovery of 465 bacterial isolates of *F. tularensis* subsp. *holarctica* from more than 27 species (Figs. 1 and S1). Initial surveillance efforts were driven by outbreaks of tularemia; our data also showed that bacterial isolates were routinely recovered in the absence of human reporting. These findings support evidence of long-standing natural foci of *F. tularensis* subsp. *holarctica* circulating among vectors and mammals in Georgia.

High numbers of bacterial isolates from rodents and vectors suggest they play a role in maintaining an enzootic cycle in Georgia, similar to findings from the former Soviet republics and Europe (Olsuf'ev and Rudnev 1960, Gyuranecz et al. 2012, Rossow et al. 2014). The common vole has been implicated in the spillover to other species and in the enzootic maintenance of the bacteria (World Health Organization

2007, Rossow et al. 2014, Hestvik et al. 2015). A study in Ukraine also identified long-standing foci of *F. tularensis* associated with the common vole and *Dermacentor* spp., among other species (Hightower et al. 2014).

In keeping with previous research, we identified repeated isolations of *F. tularensis* subsp. *holarctica* from the common vole and from blood feeding arthropods. Research in Hungary hypothesized that population dynamics of the common vole (2- to 3-year cycles) and peak densities drive the ecology and spillover of *F. tularensis* subsp. *holarctica* (Gyuranecz et al. 2012). We documented perturbations in the population density of the common vole, limited to one region, but did not find a correlation with isolation rates (Fig. S1). The common vole was the most abundant rodent trapped and had the highest rates of bacterial isolation in our study; however, the susceptibility of different mammalian species to *F. tularensis* may have influenced these findings (World Health Organization 2007). The role of the common vole in facilitating spillover to other species in Georgia through contamination of the environment requires further investigation (World Health Organization 2007, Gyuranecz et al. 2012, Rossow et al. 2014).

Interestingly, we identified several years of quiescence (no isolation) despite trapping and collecting several thousand specimens. This observation may be indicative of interepizootic periods. Recent research in Hungary suggested a European brown hare (*Lepus europaeus*)–tick cycle maintained *F. tularensis* during interepizootics (Gyuranecz et al. 2011). Although no testing of the European brown hare was conducted in our study, future surveys should focus on efforts to trap and test this species in Georgia.

Our findings provide evidence of long-standing foci of *F. tularensis* subsp. *holarctica* that may be maintained by a common vole–tick cycle. We used a more stringent method of determining positives via bacterial culturing, whereas using serology may have elicited a higher number of positive samples. Isolation rates of specimens and population dynamics of the common vole were only available for subsets of the study period, and more complete records in the future will provide a better understanding of the dynamics of *F. tularensis*. Given the history of tularemia re-emergence in Georgia, continued monitoring of small mammals, vectors, and environmental sampling may provide an early indication of outbreak risk in humans.

Acknowledgments

This work was funded by the US Defense Threat Reduction Agency's Cooperative Biological Engagement Program in Georgia through the GG-19 Project. J.K. Blackburn and I.T. Kracalik were supported through the DTRA Academic Engagement Program administered by Penn State University. This manuscript is dedicated to the memory of Dr. Ioseb Velijanashvili.

Author Disclosure Statement

No competing financial interests exist.

References

Chanturia G, Birdsell DN, Kekelidze M, Zhgenti E, et al. Phylogeography of *Francisella tularensis* subspecies *holarctica* from the country of Georgia. *BMC Microbiol* 2011; 11:139.

- Chitadze N, Kuchuloria T, Clark D, Tsertsvadze E, et al. Waterborne outbreak of oropharyngeal and glandular tularemia in Georgia: Investigation and follow-up. *Infection* 2009; 37: 514–521.
- Gyuranecz M, Rigó K, Dán Á, Földvári G, et al. Investigation of the ecology of *Francisella tularensis* during an interepizootic period. *Vector Borne Zoonotic Dis* 2011;11:1031–1035.
- Gyuranecz M, Reiczigel J, Krisztalovics K, Monse L, et al. Factors influencing emergence of tularemia, Hungary, 1984–2010. *Emerg Infect Dis* 2012; 18:1379.
- Hestvik G, Warns-Petit E, Smith L, Fox N, et al. The status of tularemia in Europe in a one-health context: A review. *Epidemiol Infect* 2015: 143:2137–2160.
- Hightower J, Kracalik IT, Vydayko N, Goodin D, et al. Historical distribution and host–vector diversity of *Francisella tularensis*, the causative agent of tularemia, in Ukraine. *Parasites Vectors* 2014; 7:453.
- Olsuf'ev N, Rudnev G. Tularemia. Moscow, 1960 1960; Translated into English.
- Petersen JM, Schriefer ME. Tularemia: Emergence/re-emergence. *Vet Res* 2005; 36:455–467.
- Rossow H, Sissonen S, Koskela KA, Kinnunen PM, et al. Detection of *Francisella tularensis* in voles in Finland. *Vector Borne Zoonotic Dis* 2014; 14:193–198.
- Velijanashvili I. Tularemia on Meskhet-Javakheti Plateau. Dissertation (in Russian), 1992.
- World Health Organization. *World Health Organization Guidelines on Tularemia*. Geneva: World Health Organization, 2007:125.

Address correspondence to:

Jason K. Blackburn

Spatial Epidemiology & Ecology Research Laboratory

Department of Geography

University of Florida

3141 Turlington Hall

Gainesville, FL 32611

E-mail: jkblackburn@ufl.edu